

JOURNAL OF ALGEBRA 47, 223-230 (1977)

On the Solvability of Certain Factorizable Groups

DAVID FINKEL

*Department of Mathematics, Bucknell University, Lewisburg, Pennsylvania 17837**Communicated by B. Huppert*

Received May 7, 1975

1. INTRODUCTION

Let G be a group and $G = AB$, where A and B are subgroups of G . There are a number of results which deduce the solvability of G from suitable conditions on A and B . In particular, in several of these results the conditions are that A and B each contain a subgroup of index 2 of particular form: for example, if G is finite and A and B are both dihedral or dicyclic, then G is solvable [2, 6]. Also see recent work of Knop [5]. We prove a result of this type, namely

MAIN THEOREM. *Let G be a finite group with subgroups A , B , H , and K such that $G = AB$, $H \leq A$, $K \leq B$, $[A : H] = [B : K] = 2$ and H is nilpotent and K is a p -group for some prime p . Then G is solvable.*

2. PRELIMINARIES

All groups considered are finite. The notations are standard. In particular, $X \leq Y$ means X is a subgroup of Y with $X < Y$ and $X \trianglelefteq Y$ meaning proper and normal subgroups, respectively, and if X is any subset of G , then $\langle X \rangle$ means the subgroup of G generated by the elements of X . Also $\pi(G)$ is the set of primes dividing the order of G . Many of the results we quote can be found in [7, especially Chap. 13].

We use throughout the Kegel-Wielandt theorem [4, 9] that the product of two finite nilpotent groups is solvable.

In addition, we need a lemma, which is essentially the proof of Kegel's result. We follow the proof given in [7, p. 381].

LEMMA. *Let G be a finite group, A and B subgroups of G with $G = AB$ and suppose every proper quotient group of G , and every proper subgroup of G containing A or B is solvable. If there is a prime p such that $p \nmid |A|$ and $p \nmid |B|$*

and if there are Sylow p -subgroups P and Q of A and B , normal in A and B , respectively, then G is solvable.

Proof. Let $N = P^G$, the normal closure of P in G . If $AN < G$, then AN is solvable, since it is a proper subgroup containing A , so N is solvable as well. Since G/N is solvable by the hypotheses of the lemma, this makes G solvable. Thus we may assume $AN = G$ and similarly $BN = G$, and $G/N = AN/N = BN/N$ has Sylow p -subgroups PN/N and QN/N . Since $P \leq N$, however, we have $PN = N$ and so $QN = N$. Thus $Q \leq N$, so $Q^G \leq N = P^G$. By symmetry we also have $P^G \leq Q^G$, so $P^G = Q^G$.

Since P and Q are the unique Sylow p -subgroups of A and B , respectively, it follows [8, Satz 6] that PQ is a subgroup of G , and from the normality of P and Q in A and B , respectively, it follows by [9, Lemma 10] that PQ^t is a subgroup for all t in G . Let P_1 be a p -subgroup of N containing P maximal with respect to the property that P_1Q^t is a subgroup of G for all $t \in G$. Then $P_1^G = N$.

If P_1 is normal in N then N is generated by p -subgroups normal in N , making N itself a p -group and therefore solvable. Since G/N is solvable by hypothesis, G is solvable in this case.

Thus we may assume that P_1 is not normal in N . Since $Q^G = N$, there exist $x, y \in G$ such that $y \in Q^x$ and $y \notin N(P_1)$. Since P_1Q^t is a subgroup for all $t \in G$, also $P_1^yQ^t$ is a subgroup, since $P_1^yQ^t = (P_1Q^{ty^{-1}})^y$. Therefore $\langle P_1, P_1^y \rangle Q^t$ is a subgroup for all $t \in G$. But from the choice of y , $P_1 < \langle P_1, P_1^y \rangle \leq N$, so from the maximality of P_1 , we conclude $\langle P_1, P_1^y \rangle$ is not a p -group.

However, $\langle P_1, P_1^y \rangle \leq \langle P_1, P_1^y \rangle Q^x = \langle P_1, Q^x \rangle = P_1Q^x$, which is a p -group. This contradiction completes the proof of the lemma.

We now prove the first of two preliminary results.

THEOREM 1. *If G is a finite group, A, B , and H subgroups of G such that $G = AB$, $H \leq A$ and $[A : H] = 2$, H is nilpotent and B is a p -group for some prime p , then G is solvable.*

Proof. Suppose G is a counterexample of minimal order. If N is any non-trivial normal subgroup of G , $G/N = AN/N \cdot BN/N$, and BN/N is a p -group and AN/N contains a subgroup HN/N of index 1 or 2, so either AN/N is nilpotent or has a nilpotent subgroup of index 2. Then G/N is solvable by the Kegel-Wielandt theorem or by induction. If M is any proper subgroup of G with $A \leq M$, then by the Dedekind identity $M = A \cdot (M \cap B)$ and so M is solvable by induction. Similarly, any proper subgroup of G containing B is solvable by the Kegel-Wielandt theorem or induction.

For any prime q dividing $|H|$, let H_q be the Sylow q -subgroup of H . Since H_q is a characteristic subgroup of H , which is normal in A , we have $H_q \triangleleft A$. If $q \neq 2$, H_q is actually a normal Sylow q -subgroup of A . Thus A has a normal 2-complement, H_2' .

We consider several cases, depending on the prime p .

Case 1. $p = 2$. Consider $G_2 \cdot H_2'$, where G_2 is a Sylow 2-subgroup of G . Since each prime of $|G|$ divides $|G_2 \cdot H_2'|$ to its full power in $|G|$, we must have $G = G_2 \cdot H_2'$, and G is solvable by the Kegel-Wielandt theorem.

Case 2. $p \nmid |A|$ and $p \neq 2$. All the conditions of the lemma are satisfied, and so G is solvable in this case also.

Case 3. $p \nmid |A|$ and $p \neq 2$. (i) G must be non-Abelian simple. For N any nontrivial normal subgroup of G , consider AN which is a subgroup of G . If $AN < G$, then AN is solvable as $A \leq AN$, and, therefore, N is solvable as $N \leq AN$. Since G/N is also solvable, this would imply that G is solvable. So $AN = G$. Similarly $BN = G$. Since in this case $|A|$ and $|B|$ are relatively prime, we have $|G| = |A| \cdot |B|$, and so $|B| \nmid |N|$ and $|A| \nmid |N|$, which imply $|G| \nmid |N|$ and $G = N$. Thus G is simple.

(ii) We must have $2 \nmid |H|$. If not, the Sylow 2-subgroups of G have order 2, and so G cannot be non-Abelian simple, contrary to (i).

(iii) Let A_2 be a Sylow 2-subgroup of A . Since $H_2 \triangleleft A_2$, we have $H_2 \cap Z(A_2) > 1$. But if $1 \neq x \in H_2 \cap Z(A_2)$, then x commutes with H_2' , since H is nilpotent, and with A_2 , so it commutes with $A = A_2 \cdot H_2'$. Thus $x \in Z(A)$. Then the conjugacy class of x in G has p -power size, since $|A| \nmid |C_G(x)|$, and so by a theorem of Burnside, G cannot be simple, contrary to (i). This completes the proof of the theorem.

We now reverse the roles of H and B .

THEOREM 2. *If G is a finite group, A , B , and H subgroups of G such that $G = AB$, $H \leq A$ and $[A : H] = 2$, H is a p -group for some prime p , and B is nilpotent, then G is solvable.*

Proof. Let G be a counterexample of minimal order. As in Theorem 1, every proper quotient group of G and every proper subgroup of G containing A or B is solvable. We now consider:

Case 1. $p = 2$. If $p = 2$, then A is a 2-group and so G is solvable by the Kegel-Wielandt theorem.

Case 2. $p \nmid |B|$. By the lemma, $p \nmid |B|$ implies that G is solvable.

Case 3. $p \nmid |B|$ and $p \neq 2$. We must have $2 \nmid |B|$. If $2 \nmid |B|$, then $|A|$ and $|B|$ are relatively prime. As in Theorem 1, this implies that G is non-Abelian simple. But since in this case a Sylow 2-subgroup of G has order 2, G is not non-Abelian simple, and so $2 \nmid |B|$.

Now, let $K = N_G(B_2)$, where B_2 is the Sylow 2-subgroup of B . If $K = G$, then B_2 is a solvable normal subgroup of G , making G solvable. Thus $K \neq G$.

Now $B \leq K$ since B is nilpotent, and so K is solvable. Let C be a Hall $\pi(B)$ -subgroup of K containing B . Now if $A \cap B > 1$, we would have $|A \cap B| = 2$ and then $G = HB$ and G would be solvable by the Kegel-Wielandt theorem. Thus $A \cap B = 1$. Therefore, if G_2 is a Sylow 2-subgroup of G containing B_2 , we have $[G_2 : B_2] = 2$, and so K contains a full Sylow 2-subgroup of G and $2 \mid |B_2| \mid |C|$. Also $|B| \mid |C|$, and since $p \nmid |B|$, we have $p \nmid |C|$. Therefore $|C| = 2 \mid |B|$. By calculating orders, we have $G = CH$. Now since H is a p -group and C has a nilpotent subgroup of index 2, G is solvable by Theorem 1, and the proof of the theorem is complete.

The simple group of order 168, $\text{PSL}(2, 7)$, is the product of a group of order 21 and a group of order 8. Thus in Theorems 1 and 2 the condition $[A : H] = 2$ cannot be replaced by $[A : H]$ is any prime. However, the proofs are valid, essentially unchanged, if this condition is replaced by $[A : H] = q$, where q is the smallest prime dividing the order of G . (Of course, then G would be solvable for having odd order.) In the proof of the main theorem, however, the prime 2 plays a critical role.

3. PROOF OF MAIN THEOREM

Let G be a counterexample of minimal order. By the Kegel-Wielandt theorem, or by Theorem 1, Theorem 2, or by induction (briefly "by induction"), every proper quotient group of G and every proper subgroup of G containing A or B is solvable. Also G has no solvable normal subgroups. We show

(1) $p \neq 2$. If $p = 2$, then B is a 2-group and so G is solvable by Theorem 1.

(2) $p \nmid |H|$. If $p \mid |H|$, let P be a Sylow p -subgroup of H . Since H is nilpotent, P is a characteristic subgroup of H , which is normal in A , so P is a normal p -subgroup of A . Also, K is a normal Sylow p -subgroup of B . The group G now satisfies all the conditions of Kegel's lemma, and so G is solvable, contradicting the choice of G .

(3) $(|A|, |B|) = 2$. This follows from (2) and the fact that $|A| = 2 \mid |H|$ and $|B| = 2 \mid |K|$.

(4) $A \cap B = 1$ and $|G| = |A| \cdot |B|$. If $A \cap B > 1$, then by (3), $A \cap B$ has order 2, and is a Sylow 2-subgroup of B , say $A \cap B = B_2$. Then $B = B_2 \cdot K$, and so $G = AB = AB_2K = AK$ and is solvable by Theorem 1. Therefore $|A \cap B| = 1$ and $|G| = |A| \cdot |B| / |A \cap B| = |A| \cdot |B|$.

(5) There exists a normal subgroup N of G , with $[G : N] = 2$.

Let \mathcal{S} be the set of right cosets of A in G , and consider the action of G on \mathcal{S} . If this action is not faithful, the kernel is a nontrivial normal subgroup of G which is contained in A and therefore solvable, making G solvable. Hence

the action is faithful. Now since $A \cap B = 1$, B acts regularly on \mathcal{S} , and \mathcal{S} has $[G : A] = |B|$ elements which is twice an odd number. Thus an involution in B is represented as an odd number of disjoint transpositions, and hence as an odd permutation. So G contains an element represented as an odd permutation, and so G has a normal subgroup of index 2, which we call N (see [6]).

Of course, since G is not solvable and any proper quotient group of G is solvable, N is not solvable.

(6) N is the only proper normal subgroup of G . Let M be any proper normal subgroup of G . Then AM is a subgroup of G , and contains A , so if $AM < G$, AM is solvable by induction, and so M is solvable, making G solvable. Hence $AM = G$. Similarly $BM = G$. From $AM = G$ and (4) follows $|B| \mid |M|$, and similarly $|A| \mid |M|$. Since $(|A|, |B|) = 2$, this implies $\frac{1}{2}|G| \mid |M|$, so $|M| = \frac{1}{2}|G|$ or $|G|$.

Now suppose $|M| = \frac{1}{2}|G|$. We wish to show $M = N$ (that is, that any two normal subgroups of this order are equal). Suppose $M \neq N$. Then since $M \cap N$ is a normal subgroup of G , $M \cap N = 1$ by the paragraph above. Then $|M| \cdot |N| = |MN| = |G| = 2|M|$, and so $|N| = 2$, and $|G| = 4$ and so G is solvable, a contradiction. Hence (6) follows.

(7) N is the direct product of isomorphic non-Abelian simple groups. Since N is the only proper normal subgroup of G , N has no proper characteristic subgroups, and N is not solvable, and so N has the claimed structure [7, Theorem 4.4.3].

We now fix some notation. By [8], there exist Sylow 2-subgroups G_2 , A_2 , and B_2 of G , A , and B , respectively, such that $G_2 = A_2B_2$. Since N is normal in G , there exists a Sylow 2-subgroup of N , N_2 contained in G_2 . Also, let H_2' be the unique Hall $2'$ -subgroup of H , and let H_2 be the Sylow 2-subgroup of H . Note that H_2 and H_2' are normal in A .

(8) $H_2' \leq N$, $K \leq N$, $A_2 \not\leq N$, $B_2 \not\leq N$. The first two just follow from the fact that N is a normal subgroup of G of index 2. If A_2 or B_2 were contained in N , then A or B would be contained in N , and by induction N would be a solvable normal subgroup of G , a contradiction.

(9) $2 \mid |H|$, since otherwise $|G_2| = 4$, and $|N_2| = 2$, which implies that N is actually simple (and not a direct product); but a non-Abelian simple group cannot have a Sylow 2-subgroup of order 2. Recall that we have $[G_2 : A_2] = [A_2 : H_2] = 2$.

(10) The following cannot occur: N simple, N_2 dihedral of order 4 or 8, and $H_2N_2 = G_2$. Assume to the contrary that this does occur. Note this implies $HN = G$, and so if we call $|N| = 2^\alpha p^\beta n$, where $\alpha = 2$ or 3 , and $(2, n) = (p, n) = 1$, then we can calculate $|H \cap N| = 2^{\alpha-2}n$ and so $H \cap N$ is a nilpotent subgroup of N of index $4p^\beta$.

Now by Gorenstein and Walter's classification of all simple groups with dihedral Sylow 2-subgroup [1], we must have N isomorphic to $\text{PSL}(2, q^f)$, $q^f > 3$, q a prime, or to A_7 . However, none of these is possible. In A_7 , a subgroup satisfying the conditions on $H \cap N$ would be Abelian; A_7 has no such subgroup. In $\text{PSL}(2, q^f)$, we can use Dickson's determination of the subgroups of $\text{PSL}(2, q^f)$ [3, Theorem II.827], and see that none of these groups has a subgroup satisfying the conditions on $H \cap N$, except $\text{PSL}(2, 4)$ and $\text{PSL}(2, 5)$, with $|H \cap N| = 3$ or 5. Of course $\text{PSL}(2, 4) \cong \text{PSL}(2, 5) \cong A_5$, the alternating group on five letters. But if $N \cong A_5$, then $G \cong S_5$, the symmetric group on five letters, as this is the only group of order 120 having a subgroup isomorphic to A_5 as its only normal subgroup. However, S_5 has no factorization of the type in the hypotheses in this theorem. This establishes (10).

(11) $H_2 \cap N_2 > 1$. If $H_2 \cap N_2 = 1$, then $2 |N_2| = |G_2| = |H_2 N_2| = |H_2| \cdot |N_2|$ and so $|H_2| = 2$. Therefore, $|G_2| = 8$ and $|N_2| = 4$. Then N cannot be the direct product of more than one non-Abelian simple group, since each would have a Sylow 2-subgroup of order 2. Hence N must be simple. Then N_2 cannot be cyclic, as a non-Abelian simple group cannot have a cyclic Sylow 2-subgroup [7, Theorem 6.2.11]. Thus N_2 is dihedral of order 4 and $H_2 N_2 = G_2$, contrary to (10). This establishes (11).

(12) There is a subgroup C , with $1 < C \leq H_2 \cap N_2$, and C normal in N_2 .

If H_2 is normal in G_2 , then $H_2 \cap N_2$ is normal in G_2 , and since $H_2 \cap N_2 > 1$ by (11), we have $C = H_2 \cap N_2$ as required.

If H_2 is not normal in G_2 , then $N_{G_2}(H_2) = A_2$, and since $[G_2 : A_2] = 2$, H_2 has exactly two conjugates in G_2 , say H_2 and H_2^x , for any $x \in G_2 - A_2$. Also, since $A_2 \triangleleft G_2$, we have $H_2^x \leq A_2$, so $H_2 H_2^x = A_2$ and

$$2 |H_2| = |A_2| = (|H_2| |H_2^x|) / (|H_2 \cap H_2^x|),$$

which implies $|H_2| = 2 |H_2 \cap H_2^x|$ and so $[H_2 : H_2 \cap H_2^x] = 2$.

Note that $H_2 \cap H_2^x$ is a normal subgroup of G_2 , as it is the intersection of all the conjugates of H_2 in G_2 .

We now consider several cases, depending on the order of $H_2 \cap H_2^x$:

(i) $|H_2 \cap H_2^x| = 1$. In this case $|H_2| = 2$ so $|N_2| = 4$. Then N cannot be the direct product of more than one non-Abelian simple group, since each would have Sylow 2-subgroup of order 2, so N must be simple. Since N_2 cannot be cyclic, as above, N_2 must be dihedral. Since $N_2 \triangleleft G_2$, we have $H_2 N_2$ a subgroup of G_2 , and either $H_2 N_2 = N_2$ or $H_2 N_2 = G_2$. The latter cannot occur by (10). If $H_2 N_2 = N_2$, then $H_2 \cap N_2$ has order 2 and $H_2 \cap N_2 \triangleleft N_2$ since it has index 2; thus in this case we let $C = H_2 \cap N_2$.

(ii) $|H_2 \cap H_2^x| = 2$. In this case $|N_2| = 8$ and, as in (i), N must be simple. Again $H_2N_2 = N_2$ or $H_2N_2 = G_2$. If $H_2N_2 = N_2$, then $H_2 = H_2 \cap N_2$, and $H_2 \cap H_2^x \leq H_2 \cap N_2$, and $C = H_2 \cap H_2^x$ is as required in this case.

If $H_2N_2 = G_2$, then $|H_2 : H_2 \cap N_2| = 2$, so $|H_2 \cap N_2| = 2$. Since N_2 has order 8, N_2 is either Abelian, quaternion, or dihedral. N_2 cannot be dihedral, by (10). If N_2 is either Abelian or quaternion, the subgroup $H_2 \cap N_2$ of order 2 is normal in N_2 , and so $C = H_2 \cap N_2$ is as required in this case.

(iii) The remaining case is $|H_2 \cap H_2^x| \geq 4$. Since $H_2 \cap H_2^x \leq G_2$ and $|G_2 : N_2| = 2$ we have $|H_2 \cap H_2^x \cap N_2| \geq 2$, so $C = H_2 \cap H_2^x \cap N_2$ is as required (in fact, $C \triangleleft G_2$). Thus N_2 has a normal subgroup C , as claimed.

(13) Conclusion. Since $1 < C \triangleleft N_2$, we have $1 < C \cap Z(N_2)$. Let $x \in C \cap Z(N_2)$, $x \neq 1$. Then $N_2 \leq C_N(x)$; since $x \in H_2$ and H is nilpotent, $H_2 \leq C_N(x)$. Thus $[N : C_N(x)]$ is a power of p , since all the other primes of N divide $C_N(x)$ to their full power in N , and so the size of the conjugacy class of x in N is a power of p . Thus N cannot be non-Abelian simple by the theorem of Burnside quoted above.

Thus N must be a direct product of non-Abelian simple groups; let N^* be one of those non-Abelian simple groups for which the projection of x is not the identity. Call the projection $\pi : N \rightarrow N^*$. Then $C_{N^*}(\pi(x)) = \pi(C_N(x))$. Since $N_2 \leq C_N(x)$, we have that $\pi(N_2)$, which is a Sylow 2-subgroup of N^* , is contained in $C_{N^*}(\pi(x))$. Similarly $C_{N^*}(\pi(x))$ contains Sylow subgroups of N^* for all primes of N^* different from p . Therefore, $[N^* : C_{N^*}(\pi(x))]$ is a power of p , which by Burnside's theorem, contradicts N^* being non-Abelian simple.

This final contradiction completes the proof of the Main Theorem.

ACKNOWLEDGMENTS

The author wishes to thank Research Corporation, which supported this work through a Cottrell College Science grant. He also wishes to thank Terri K. Bourdon, a participant in that grant, and Professor William R. Scott, who made helpful comments on an earlier version of this work.

REFERENCES

1. D. GORENSTEIN AND J. H. WALTER, The characterization of finite groups with dihedral Sylow 2-subgroups, I, II, III, *J. Algebra* **2** (1965), 85–151, 218–270, 334–393.
2. B. HUPPERT, Über die Auflösbarkeit factorisierbarer Gruppen, *Math. Z.* **59** (1953), 1–7.
3. B. HUPPERT, "Endliche Gruppen I," Springer-Verlag, Berlin, 1967.
4. O. KEGEL, Producte nilpotenter Gruppen, *Arch. Math.* **12** (1961), 90–93.
5. L. E. KNOP, Sufficient conditions for the solvability of factorizable groups, *J. Algebra* **38** (1976), 136–145.

6. W. R. SCOTT, Solvable factorizable groups, *Ill. J. Math.* 1 (1957), 389-394.
7. W. R. SCOTT, "Group Theory," Prentice-Hall, Englewood Cliffs, N.J., 1964.
8. H. WIELANDT, Über das Product paarweise vertauschbarer nilpotenter Gruppen, *Math. Z.* 55 (1951), 1-7.
9. H. WIELANDT, Über Producte von nilpotentent Gruppen, *Ill. J. Math.* 2 (1958), 611-618.